

HARMFUL • ECOTOXIC • CORROSIVE

KEEP OUT OF REACH OF CHILDREN



Nufarm

TRIMEC®

FREE RECYCLING




AGRECOVERY

www.agrecovery.co.nz

Active Ingredients: 600g/litre mecoprop, 150g/litre MCPA and 18.7g/litre dicamba in the form of a soluble concentrate.

A selective herbicide for the control of a wide range of broadleaf weeds in wheat, barley, oats and ryegrass seed crops

Net: 20 Litres



9 400588 033178 >

04-18/PS/A0583-20LNZ1.2



Nufarm

Grow a better tomorrow.

HAZARD CLASSIFICATION

6.1D, 6.3B, 6.5A, 6.5B, 6.8B, 6.9A, 8.3A, 9.1A, 9.2A, 9.3C

APPROVED HANDLER

This product must be under the control of an approved handler during use.
This product must not be used for any purpose, or in any manner, contrary to this label unless authorised under appropriate legislation.

TOXICITY

- Harmful if swallowed, inhaled or absorbed through the skin. May cause skin sensitisation or mild skin irritation from prolonged skin contact. May cause reproductive/development damage from repeated oral exposure.
- Toxic - may cause respiratory sensitisation. May cause liver or kidney damage from repeated oral or inhalation exposure at high doses.

CORROSIVE

- Danger - this product is corrosive and may cause eye damage.

ECOTOXIC

- Very toxic to aquatic organisms with long-lasting effects. Avoid contamination of any water supply with product or empty container.
 - Very toxic to soil environment. Harmful to terrestrial vertebrates.
 - Selective herbicide - very toxic to some plant species. Do not apply this product on or near desirable trees or plants or in locations where the chemical may be washed or moved into contact with their roots.
- Beware: Spray drift hazard. Apply this product carefully. Spray drift may cause serious damage to other desirable plants.** Ensure compliance with relevant regional air management plans.

HANDLING PRECAUTIONS

STORAGE

- Store in original container tightly closed and in a locked, dry, cool area away from foodstuffs, seeds and fertilizers. Storage must be in accordance with NZS 8409 Management of Agrichemicals.
- Stores containing 100L of Trimec require bunding and are subject to signage and more than 100L require emergency response plans.

PERSONAL PROTECTION

- Avoid skin or eye contact and inhalation of spray mist.
- When handling wear eye protection, waterproof gloves, boots and overalls. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.

EQUIPMENT:

- Apply using standard ground and aerial spraying equipment.

DISPOSAL

- Triple rinse empty container and add rinsate to spray tank. Recycle empty container. Otherwise crush and bury in a suitable landfill.
- Dispose of product only by using according to the label, or at an appropriate landfill.

First Aid: If swallowed do NOT induce vomiting. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. DO NOT SCRUB THE SKIN. If the concentrate is splashed in the eye, flush with running water immediately. If inhaled move the victim to fresh air immediately. Begin artificial respiration if breathing has stopped.

Spillage: Wear appropriate protective clothing and prevent material from entering waterways. Contain liquid spills and absorb with inert material. Dispose of waste safely in an approved landfill.

Transport: Do not carry this product on a passenger service vehicle.

Record Keeping: Records, as described in the current NZS 8409 Management of Agrichemicals, must be kept whenever TRIMEC is used.

Resistance Management: Trimec contains **GROUP 01 HERBICIDES** (Mecoprop and MCPA) belonging to the phenoxy group of chemistry and a **GROUP 02 HERBICIDE** (Dicamba) belonging to benzoic acid group of chemistry. Resistance to these herbicides could develop from excessive use. To minimise this risk use strictly in accordance with label instructions and resistance management strategies (www.resistance.nzpps.org/) if any for the crops listed on this label.

Mixing: TRIMEC mixes readily with water, and the use of additional wetting agent is not recommended. Pour the measured quantity of TRIMEC into the partly filled spray tank, then top up with water and agitate continually during filling.

Cleaning: Flush equipment thoroughly with clean water after use. If the spray unit which has applied TRIMEC is to be used for the application of other chemicals to sensitive crops, ensure the unit is completely decontaminated.

GENERAL INFORMATION

TRIMEC controls a wide range of broadleaf weeds including many difficult to control weeds of particular importance in the intensive cereal cropping districts. TRIMEC is particularly selective to varieties of wheat, barley and oats once these plants have reached the 4 leaf growth stage, and is suitable for use on ryegrass seedcrops once the ryegrass has developed to the 4-6 leaf stage of growth. It provides effective control of wireweed, willow weed, Californian thistle, spurrey, stinking mayweed and chamomiles as well as most other common broadleaf weeds of cereals and ryegrass seed crops.

WEEDS CONTROLLED:

1. Seedling weeds effectively controlled with TRIMEC at 3 to 3.5 litres per hectare:

Black nightshade, mouse-ear chickweed, catchfly, cornbind, docks, fathen, hedge mustard, mallow, redroot, spurrey, shepherd's purse, thistles (noddling, Scotch, sow and winged, as well as Californian thistle regrowth from rootchips), twincress, wild turnip, willow weed and wireweed.

2. Seedling weeds effectively controlled with TRIMEC at 3.5 to 4 litres per hectare:

Chamomiles, cleavers, fumitory, scrambling speedwell, stinking mayweed, water pepper and yarrow.

DIRECTIONS FOR USE:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice : Maximum Residue Levels for Agricultural Compounds.

WHEAT, BARLEY, OATS, RYEGRASS seedcrops - broadleaf weeds:

Rate of Application:

Apply 3 to 4 litres TRIMEC in 150 (minimum) to 200 litres of water per hectare. The rate to be used will be influenced by the susceptibility of the particular weeds present in the cereal crop, and their stage of growth. Early recognition of the weed problem is important so that TRIMEC may be applied when the weeds are young and making active growth. Use the 3.5 litres per hectare rate when seedling weeds have advanced to the 6-8 true leaf stage. Use the 4 litres per hectare rate where chamomiles, stinking mayweed, yarrow and water pepper are predominant.

Time of Application:

In wheat, barley and oats, TRIMEC may be applied at the recommended rate as soon as the cereal plants have reached the 4 leaf growth stage.

In ryegrass seedcrops TRIMEC should be applied over the late autumn-early winter period when the ryegrass has reached at least the 4-6 leaf stage of growth.

TRIMEC should not be applied after the cereal or ryegrass crop has reached the 'jointing' or 'node' stage.

Caution:

As TRIMEC is very damaging to clovers do not apply to crops undersown with clovers. Do not apply TRIMEC to any crop from which the straw may be required for horticultural mulching. Do not apply TRIMEC if rain is imminent, and 18 - 24 hours fine weather following is desirable. Do not exceed recommended dosage rate. When spraying do not overlap or 'spray out' corners. Poisonous plants may become palatable after spraying with this material and stock should be kept away from the area until such plants have died down.

© TRIMEC is the registered trademark of Nufarm Technologies USA

Registered pursuant to the ACVM Act 1997, No. P3253. See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to the HSN0 Act 1996, Approval Code HSR000348. See www.epa.govt.nz for approval controls.

CONDITIONS OF SALE:

CAUTION: READ THIS NOTICE BEFORE OPENING THE CONTAINER

If you open this container you will be deemed to have accepted the conditions below.

The results from use of this product may be affected by factors beyond Nufarm's control, including mixing, use, climatic conditions, time of application, crop or crop stage and the possible development of resistance to the active ingredient, even though the product is reasonably fit for the purposes stated in this label, and even though label directions are followed. Nufarm will replace, free of charge, any product which is found to have a manufacturing defect. To the extent permitted by law, all other warranties, guarantees, representations and conditions, whether expressed or implied by law, trade, custom or otherwise are expressly excluded.

When you use this product, you agree that neither Nufarm Limited nor any manufacturer of any component of the product will be liable for any damage, injury or loss including consequential loss which arises in any way from the storage, handling, or use of the product. This exclusion of liability includes loss caused by negligent acts or omissions of Nufarm Limited or any supplier or manufacturer of the product or any of their employees and agents.

Marketed by: Nufarm Limited
6 Manu Street, Otahuhu, Auckland, NZ

Telephone 0-9-270 4157

www.nufarm.co.nz

Nufarm Emergency Phonenumber: 0800-651 911 toll free 24 hours

Batch No.:

D.O.M.:

PROPER SHIPPING NAME	CLASS	UN No.	HAZCHEM	PG
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (MCPA 15% and dicamba 1.87%)	9	3082	2X	III