



CONDOR[®]

For the control of a wide range of leaf and stem diseases of Wheat, Barley, Oats, Ryegrass seed crops and Turf grass and for the treatment of Prairie grass seed for Head Smut

GROUP

3

FUNGICIDE

Contains 250g/L propiconazole in the form of an emulsifiable concentrate



AGRISOURCE
PRODUCTS FOR AGRICULTURE

**WARNING
ECOTOXIC**

Keep out of the reach
of children

Net Contents

10 Litres

FREE RECYCLING



www.agrecovery.co.nz

GENERAL INFORMATION

Condor is a systemic fungicide for the control of various foliar diseases of small grain cereals and ryegrass, as well as the treatment of prairie grass seed for Head Smut.

Wheat: Stripe Rust, Leaf Rust (Brown Rust), Speckled Leaf Blotch and Glume Blotch

Barley: Net Blotch (both net and spot forms), Spot Blotch, Leaf Scald, Leaf Rust and Ovarial Leaf Spot

Oats: Crown Rust

Ryegrass: Stem Rust and Leaf Rust (Crown Rust)

Turf: Dollar Spot, Brown Patch, Red Thread, Fusarium Patch

Prairie grass (seed treatment): Head Smut

It is an offence to use this product on animals.

IMPORTANT - RESISTANCE MANAGEMENT

Condor contains a **GROUP 3 FUNGICIDE** propiconazole. Resistance to this fungicide and related products could develop from repeated use. To minimise this risk use strictly in accordance with the label instructions. In cereals use a maximum of three DMI applications per crop, including seed treatment, at full recommended label rates.

MIXING

Add the required amount of **Condor** to the partly filled spray tank and add the remainder of the water with agitation.

COMPATIBILITY

Condor is compatible with most of the commonly used pesticides, including herbicides, used on cereals.

APPLICATION

Maximum application as a seed treatment: **Condor** must not be applied at rate exceeding 15g propiconazole per 100 kg seed.

Maximum application rate for wide dispersive use: **Condor** must not be applied at rates exceeding 125g propiconazole/ha. **Condor** must not be applied to the same area more than three times in any 364 day period.

Aerial Application: Apply in sufficient water to ensure more than 20 droplets per square centimetre on all leaves. As there are many factors involved in aerial application, for optimum results, contact your local supplier representative for detailed information.

Treated Seed: Containers of treated seed must be labelled accordingly. When used as a seed treatment, **Condor** must be applied in a purpose-built seed coating machine, in a treatment plant with local exhaust ventilation. At the time of treatment, seed must be dyed so as to differentiate it from untreated seed. Treated seed must be completely covered by soil when sown. Excess treated seed must not be left in areas accessible to birds. Personal protection equipment must be used when handling seeds treated with **Condor**. Seeds treated with **Condor** must be disposed of in accordance with the disposal requirements for toxic and ecotoxic substances. Seed treated with **Condor** must not be used for human or animal consumption.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds

WITHHOLDING PERIODS

Cereals (silage and grain crops).....42 Days

Ryegrass, Prairie grass seed crops30 Days

Turf DO NOT graze or feed clippings to livestock

CONDITIONS OF SUPPLY

All goods supplied by us are produced under rigid quality control procedures which assure the purchaser that at the time of sale they are physically suitable for use in the manner stated. However, to the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, Agri Solutions Ltd as registrant, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

REGISTRANT

Agri Solutions Ltd
Level 1, 26 Crummer Road
Grey Lynn, Auckland 1021

DISTRIBUTOR

AgriSource (2000) Ltd
45 Kitchener Road, Pukekohe 2120
09 237 0442 - www.agrisource.co.nz

BATCH NO:

DATE OF
MANUFACTURE:

CROP	DISEASE	RATES	REMARKS
Wheat	Stripe Rust, Speckled Leaf Blotch	500ml in a minimum of 200 litres water/ha	Apply as a preventative spray on susceptible cultivars. For control on less susceptible cultivars, apply at first sign of the diseases. Note: DO NOT apply Condor to silage cereal crops after GS 39, flag leaf fully unrolled, to ensure the 42-day withholding period is observed.
	Glume Blotch		Apply as a preventative spray at ear emergence on susceptible cultivars especially during cool, moist conditions which favour spread of this disease to the head. On less susceptible cultivars, apply at first sign of the disease. Note: DO NOT apply Condor to silage cereal crops after GS 39, flag leaf fully unrolled, to ensure the 42-day withholding period is observed.
Wheat, Barley	Leaf Rusts	500ml in a minimum of 200 litres water/ha	Apply as a preventative spray at ear emergence on susceptible cultivars. For control on less susceptible cultivars, apply at first sign of the disease. Note: DO NOT apply Condor to silage cereal crops after GS 39, flag leaf fully unrolled, to ensure the 42-day withholding period is observed.
Barley	Net Blotch, Spot Blotch, Leaf Scald, Ovarial Leaf Spot		Apply at first sign of the diseases. Note: DO NOT apply Condor to silage cereal crops after GS 39, flag leaf fully unrolled, to ensure the 42-day withholding period is observed.
Oats	Crown Rust		Apply as a preventative spray at flag leaf emergence on susceptible cultivars. On less susceptible cultivars, apply at first sign of the disease up to the completion of flowering. Note: DO NOT apply Condor to silage cereal crops after GS 39, flag leaf fully unrolled, to ensure the 42-day withholding period is observed.
Ryegrass seed crops	Stem and Leaf Rusts (Crown Rust)		For optimum control on susceptible cultivars, apply Condor as a preventative spray after closing up for seed, and repeat at 3 to 4 week intervals depending upon continuing disease pressure. For control on less susceptible cultivars, apply at first sign of disease.
Turf Grass	Apply in a minimum of 10 litres of water/100 square metres. Apply as a preventative spray before the signs of disease appear, when conditions are favourable for disease development. For maximum control of turf diseases, maintain the appropriate water and fertiliser requirements to minimise disease development. DO NOT mow within 24 hours of application, and delay application to after the morning dew has dried, to minimise the spread of diseases. Note: Some reduction in grass growth may occur. DO NOT apply to newly sown turf.		
	Dollar Spot	20ml/100 square metres	Apply every 14 to 28 days.
	Brown Patch	30ml/100 square metres	Apply every 10 to 21 days. Under conditions of high temperature and humidity, repeat application at the shorter interval.
	Red Thread		Apply every 14 to 21 days.
	Fusarium Patch		Make the first application in late autumn and repeat as necessary, ensuring no more than 3 applications per season.
Prairie grass seed	Head Smut	60ml/100 kg of dressed seed	Apply as a seed coat in sufficient water to ensure adequate coverage. At the time of treatment, the seed must be dyed a distinctive colour (see under 'Treated Seed').

® Condor is a registered trademark
of Agri Solutions Limited

Registered pursuant to the ACVM Act 1997 No. P009817
See www.foodsafety.govt.nz for registration conditions
Approved pursuant to HSN0 Act 1996 No. HSR101041
See www.epa.govt.nz for approval conditions

READ LABEL COMPLETELY BEFORE USE

HSNO CLASSIFICATIONS: 3.1D, 6.1E, 6.3B, 6.4A, 6.9B, 9.1A, 9.3C

RECORD KEEPING

Where 3 kg or more of the substance is applied within 24 hours, in a place where the substance is likely to enter air or water and leave the application area, a PCBU with management or control of the substance must ensure that a written record is kept of each application of the substance. Refer to Part 4 Clause 48 of the Hazardous Substances (Hazardous Property Controls) notice 2017 for details of information that must be recorded.

PCBU RESPONSIBILITIES

A PCBU with management and control of the place to where this product is being used, handled, manufactured, stored or applied must ensure that every worker who handles it, is provided with adequate information, training and instruction. This includes provision and training in the use of any personal protective equipment to be used.

QUALIFICATIONS REQUIRED FOR USE OF THIS CLASS 9 PRODUCT

Anyone who mixes, loads, or applies this product, either by air or ground based methods, is required to be suitably qualified. Refer to Schedule 10 of the Hazardous Substances (Hazardous Property Controls) notice 2017 for a list of suitable qualifications.



WARNING

Combustible liquid. May be harmful if swallowed. Causes mild skin irritation. Causes eye irritation. May cause damage to Liver through prolonged or repeated exposure.

ECOTOXIC

Very toxic to aquatic organisms. Harmful to terrestrial vertebrates. **Condor** must not be applied into or onto water. Avoid contamination of any water supply with chemical or empty container.

HANDLING PRECAUTIONS

Personal Protection: When mixing or applying, avoid contact with skin and eyes. Wear full protective clothing, chemical resistant PVC coated gloves and goggles. Do not eat or drink while using. Wash hands and face before meals and after work. Wash protective clothing after work. **Keep away from heat/sparks/open flames and hot surfaces. No smoking. Do not breathe mist, vapours or spray.**

Equipment: Apply using accurately calibrated and maintained equipment in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS8409). Flush spray equipment thoroughly with water after use.

Storage: Store in original container tightly closed and in a locked, dry, cool, well ventilated area away from fertilisers, feed, seeds and foodstuffs. Avoid temperatures below minus 5°C or above 35°C. Keep out of reach of children. Store in accordance with NZ 8409 Management of Agrichemicals.

Shelf Life: When stored appropriately this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

Spillage: Wear suitable protective clothing including face shield, impervious gloves and boots. Prevent the product or spilled material from entering water bodies. Absorb liquid spills with inert material such as earth or sand and place in waste containers. Wash area with water and absorb with further inert material. Dispose of safely.

Product Disposal: Dispose of this product only by using according to the label, or at an approved landfill or other approved facility.

Container Disposal: Triple rinse container and add rinsate to spray tank. Dispose of cleaned container at your local AGRECOVERY container collection site.

Safety Data Sheet: Additional information is provided in the SDS available from your supplier.

FIRST AID

If swallowed DO NOT induce vomiting. If medical advice is needed, have product container or label at hand. For advice, contact the National Poisons Centre 0800 Poison (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. If concentrate is splashed in the eyes, flush with running water immediately.

EMERGENCY RESPONSE

In a transport emergency call 111, Police or Fire Brigade

24 HOUR EMERGENCY RESPONSE 0800 CHEMCALL (0800 243 622)