

WARNING

KEEP OUT OF REACH OF CHILDREN



Ridomil Gold[®] MZ WG

Systemic Fungicide

syngenta[®]

Active Ingredients:

RIDOMIL Gold MZ WG contains 40 g/kg METALAXYL-M plus 640 g/kg MANCOZEB in the form of a water dispersible granule.

For the control of Early and Late Blight in Potatoes and Tomatoes, Dryberry in Canefruit, Downy Mildew in Grapes, Onions, Cucurbits and Brassica seedbeds, Phytophthora Spear Rot in Asparagus, Phytophthora Crown Rot in Asparagus transplants, and Phytophthora Root Rots in Tree nurseries.

Net Contents: 5 kg

*Registered pursuant to the ACVM Act 1997, No. P 5658.
See www.foodsafety.govt.nz/industry/acvm for registration conditions*

Approved pursuant to the HSNO Act 1996, Approval Code HSR000470.

Syngenta Crop Protection Limited

Tower 2, Level 7, 110 Symonds Street, Auckland

TM

PLEASE READ THE LABEL COMPLETELY BEFORE USE

This product MUST NOT be used for any purpose or in any manner contrary to the controls and conditions of this registration.



HSNO Classification 6.5B, 6.9B, 9.1A

WARNING

May cause sensitisation from prolonged skin contact.
May cause liver damage from repeated oral exposure at high doses.
Very toxic to aquatic organisms.

PRECAUTIONS

Store in original container, tightly closed, away from foodstuffs.
When mixing or applying wear protective clothing, including overalls, water proof gloves and footwear.
Avoid contact with skin and eyes, and inhalation of spray mist.
DO NOT eat, drink or smoke while using.
Wash hands and face before eating, drinking or smoking and after work.
Wash protective clothing daily after use.
Apply with well-maintained and calibrated spray equipment.
DO NOT re-enter site until spray has dried.
Records of use must be kept.
Application, storage and record keeping should be as described in the Management of Agrichemicals NZS8409.
DO NOT carry on a passenger service vehicle.
Avoid contamination of any water supply with chemical or empty container.

Container disposal: Ensure the container is empty.

(For Hermetic Boxes) Rinse the container and add the residue to the spray tank. If allowed under local Territorial Authority District Plans, burn if circumstances, especially wind direction permit.

(For HDPE Bottles) Triple rinse the empty container and add rinsate to the spray tank. Recycle empty container through Agrecovery (0800 247 326, www.agrecovery.co.nz). Otherwise crush and bury in an approved local authority facility. DO NOT reuse this container for any other purpose

Product disposal: Dispose of this product only by using according to the label, or at an approved landfill or other approved facility.

FIRST AID

If swallowed DO NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. Begin artificial respiration if breathing has stopped. Use mouth to nose rather than mouth to mouth. Obtain medical attention. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. DO NOT scrub the skin. If splashed in the eyes, wash out immediately with water. If inhaled, move the victim to fresh air immediately. Begin artificial respiration if breathing has stopped.

SPILLAGE

Wear suitable protective clothing including face shield, impervious gloves and boots. Prevent the product or spilled material from entering water bodies. Absorb liquid spills with inert material such as earth or sand and place in waste containers. Wash area with water and absorb with further inert material. Dispose of safely.

DISCLAIMER

Warning: RIDOMIL Gold MZ WG contains a metalaxyl isomer. Experience has shown that some strains of fungus resistant to metalaxyl isomers have developed resulting in a reduction in the efficacy of the product and thus in yield losses. Since the occurrence of resistance cannot be foreseen, Syngenta accepts no responsibility for any loss of, or damage to, crops resulting from the failure of RIDOMIL Gold MZ WG to control resistant strains. Where products have previously been found to be ineffective in controlling diseases, or if the user suspects that a tolerant or resistant strain is present, then RIDOMIL Gold MZ WG should NOT be used. Advice as to alternative treatments should be sought in such cases.

GENERAL INFORMATION

RIDOMIL Gold MZ WG is a systemic and protectant fungicide. Metalaxyl-M is systemic and is rapidly taken up by green tissue within 30 minutes and distributed within the leaves and stems providing disease control by inhibiting fungal growth within the plant. Mancozeb is a protectant, which provides a residual protective film over the plant surface, inhibiting spore germination.

RESISTANCE MANAGEMENT

Resistance to this fungicide and related products could develop from excessive use. To minimise this risk use strictly in accordance with label instructions.

DIRECTIONS FOR USE

Important: DO NOT use as an eradicant treatment (ie. once disease symptoms are visible).

Volume of Application: For broadcast spraying, use sufficient water to obtain coverage without run-off, to achieve the maximum systemic and protectant effect from the product. For high volume spraying, the rate suggested is applied to run-off. For concentrate spraying, adjust dilution rate accordingly. The addition of a surfactant is recommended with all foliar applications of RIDOMIL Gold MZ WG.

It is an offence to use this product on animals

Crop	Disease	Rate	Critical Comments
Asparagus	Phytophthora Spear Rot	12.5 to 15 kg in 200 litres water/treated hectare.	Use the higher rate if disease was evident during the previous season. Apply to weed free soil in early spring 3 to 4 weeks prior to spear emergence, either as a 50 cm wide band over the row or as a broadcast application. About 10 mm rainfall is desirable to move the fungicide into the soil.
Asparagus transplants	Phytophthora Crown Rot	500 g /130 litres of water	Immerse crowns for 5 minutes prior to transplanting.
Blackberries, Boysenberries and other Brambles	Dryberry and Canespot	250 g /100 litres water	Mature Vines: Apply at 14 day intervals from bud burst to pre-flowering, reducing to 10 day intervals from tight cluster to early fruit formation when dryberry infection can be most serious. Primocanes: Apply immediately after final harvest and repeat at 14 day intervals while warm, wet conditions favour the disease. Under dry conditions use alternative protectant fungicides according to label recommendations followed by one application of RIDOMIL Gold MZ WG before leaf fall.
Brassica Seedbeds	Downy Mildew	20 g /100 sq metres of seedbed (2 kg/ha)	Use sufficient water to obtain complete coverage. Make the first application as soon as many of the seedlings have emerged, and repeat at 10 day intervals during transplanting.
Cucurbits	Downy Mildew	2.5 kg in 500 to 700 litres water/ha	Make a maximum of 3 applications at 10 to 14 day intervals during the period of active growth when conditions of low night temperatures with dew formation favour the disease.

Crop	Disease	Rate	Critical Comments
Grapes	Downy Mildew	250 g /100 litres of water applying a minimum of 1000 litres/ha.	Make a maximum of 3 applications at 14 day intervals, when warm and wet conditions favour the disease at the pre-flowering, post-flowering, and early fruit set stages of growth. Subsequent applications of alternative protectant fungicides can be made as and when necessary until harvest. An application of RIDOMIL Gold MZ WG can be made after harvest, but before leaf fall to minimise carryover of disease into the following season.
Onions	Downy Mildew	2 kg in 500 to 700 litres water/ha	Make a maximum of 3 applications at 10 day intervals during the period of active growth when conditions of high humidity favour the disease.
Potatoes, Tomatoes	Early Blight, Late Blight	2.5 kg in 500 to 750 litres water/ha, or use 250 g/100 litres of water as a foliar spray to run-off.	Make a maximum of 3 applications at 10 to 14 day intervals during the period of active growth. Make the first application when the crop has fully emerged and before disease has occurred. Use the shorter spray interval when weather conditions favour the disease especially during active crop growth in the period prior to flowering. Apply when foliage is dry and no rain or irrigation is expected for at least 2 hours. DO NOT apply in the period following the completion of flowering as the translocation of RIDOMIL Gold MZ WG is considerably reduced if applied when haulms are drying off. IMPORTANT: Bacterial Speck and Bacterial Spot: As RIDOMIL Gold MZ WG does NOT control bacterial diseases in tomatoes, it may be necessary to use a protectant such as cupric hydroxide or copper oxychloride. Botrytis: RIDOMIL Gold MZ WG does NOT control Grey Mould (<i>Botrytis cinerea</i>).
Tree Nurseries (radiata pine, kauri, cypresses)	Phytophthora Root Rots	12.5 to 50 kg in 200 to 300 litres water/treated hectare, preferably to damp ground.	Use the higher rates for soils with high organic content, and on beds with a previous history of root rot problems. Apply to seedbeds before seedling emergence. About 10 mm rainfall or irrigation is desirable to move the fungicide into the soil.

MIXING

RIDOMIL Gold MZ WG mixes readily with water. Pre mix the product with sufficient water into a smooth cream, pour into the partly filled spray tank and add the remainder of the water with agitation ensuring complete mixing before spraying. Continue agitation during spraying and after a stoppage before recommencing spraying.

COMPATIBILITY

RIDOMIL Gold MZ WG is compatible with most of the commonly used insecticides and fungicides except lime sulphur.

WITHHOLDING PERIODS

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residue Limits of Agricultural Compounds) Food Standards.

Asparagus.....	21 days
Canefruit	14 days
Cucurbits	4 days
Grapes.....	14 days
Onions	14 days
Potatoes	14 days
Tomatoes (field),	14 days
Tomatoes (glasshouse).....	3 days

STORAGE

Store in a dry, shaded area, avoiding temperatures below minus 10°C and above 35°C. See Safety Data Sheet for additional storage information. Stores containing quantities of 100 kg or more require signage and secondary containment.

SAFETY DATA SHEET

If additional hazard or safety information is required, refer to the Safety Data Sheet. For a copy visit our website at www.syngenta.co.nz

DISCLAIMER

This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. Syngenta's liability for breach of the express or any non-excludable implied warranty is limited to product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which Syngenta shall have no liability

Product names marked ® or ™, the ALLIANCE FRAME the SYNGENTA Logo and the PURPOSE ICON are Trademarks of a Syngenta Group Company



Distributed by and registered to Syngenta Crop Protection Limited, Tower 2, Level 7, 110 Symonds Street, Auckland.

Bar-Code: 9-414644-00019 (plus check)

Batch Number:	
Date of Manufacture:	



HDPE only

UN 3077
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, SOLID, N.O.S.
(mancozeb)
MARINE POLLUTANT
PACKING GROUP III, HAZCHEM 2Y
CONTAIN SPILLAGE



In a transport emergency dial 111, Police or Fire Brigade.
For specialist advice in an emergency only, call 0800 734 607 (24 hours).