

Hussar®

Net contents : 600 g

Herbicide


Claim statement: Selective post-emergence herbicide for the control of certain broad-leaved and grass weeds in autumn and spring-sown wheat, durum wheat, triticale, ryecorn and barley

Contains: 50g /kg iodosulfuron-methyl-sodium and 150g/kg mefenpyr-diethyl(safener) in the form of a water dispersible granule

Registered pursuant to the ACVM Act 1997, No P7047
See www.foodsafety.govt.nz for registration conditions.


Approved pursuant to the HSNO Act 1996, No HSR000065
See www.epa.govt.nz for approval conditions

DG Information



UN 3082
PROPER SHIPPING NAME
PACKING GROUP I/II/III
CLASS 3/6/8/9
HAZCHEM CODE

IN A TRANSPORT EMERGENCY DIAL 111 POLICE OR FIRE BRIGADE



DANGER**HSNO Classifications:**

- 6.3B Causes mild skin irritation.
 8.3A Causes serious eye damage
 9.1A Very toxic to aquatic life with long lasting effects
 9.2A Very toxic to the soil environment

PRECAUTIONS:

- Read label before use
 Do not eat, drink or smoke while using this product.
 Remove protective clothing and wash hands and face thoroughly before meals and after work.
 Avoid contact with the skin and eyes
 Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area
 Do not apply directly into or onto water

PERSONAL PROTECTION:

When handling, wear full protective clothing such as impervious gloves, respirator, waterproof hat, coat and trousers (worn outside rubber boots).

STORAGE: Store in original container tightly closed and in a locked dry, cool, well-ventilated area, away from foodstuffs.

SPILLS AND DISPOSAL: When dealing with spills wear personal protective clothing and equipment as described in the PERSONAL PROTECTION section.

Contain the spill by damming, recover spilt product by absorbing with sawdust or an inert absorbent material then transfer the recovered spilt material to a properly labelled drum. Deal with all spillages immediately. If contamination of drains, streams, watercourses, etc. is unavoidable, warn the local water authority.

Disposal: Dispose of this product only by using according to the label or at an approved landfill. Avoid contamination of any water supply with product or empty container.

Container Disposal: Triple rinse container and add residue to mixture. Take to an Agrecovery collection site.

FIRST AID:

If swallowed, do NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin contact occurs, remove contaminated clothing and flush skin with running water. If splashed in eyes, wash out immediately with water, if irritation persists get medical advice. If inhaled, move the victim to fresh air immediately.

BEWARE: APPLY THIS PRODUCT CAREFULLY. SPRAY DRIFT MAY CAUSE SERIOUS DAMAGE TO OTHER DESIRABLE PLANTS.



Hussar is a selective post-emergence herbicide for the control of certain broad-leaved and grass weeds in wheat, durum wheat, triticale, ryecorn and barley.

DIRECTIONS FOR USE:

MIXING: Half fill the spray tank with water, then with agitation in motion, add the correct amount of **Hussar** directly into the spray tank. Add other relevant compatible pesticides and finally, add the Partner®. Complete filling the tank with agitators in motion.

Agitation must continue before and during spraying.

Surfactant: The addition of Partner is essential for control.

Application: Apply through any standard crop sprayer capable of giving uniform coverage at a spray volume of 200 - 300 l/ha. Use spray pressures of at least 200 kPa and select nozzles capable of producing fine spray droplets. Avoid overlaps.

RESISTANCE MANAGEMENT

Group	B	Herbicide
-------	----------	-----------

Hussar is a member of the sulfonyl urea group of herbicides and inhibits ALS. Some naturally occurring weed biotypes resistant to **Hussar** and other ALS herbicides may exist through normal genetic variability in any weed population. Resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. Resistant weeds will not be controlled by **Hussar** or other ALS herbicides. Since occurrence of resistant weeds is difficult to detect prior to use, Bayer accepts no liability for any losses that may result from failure of **Hussar** to control resistant weeds. Chickweed biotypes resistant to Glean® have been identified. To prevent or delay the development of resistance, use **Hussar** in tank mixes where appropriate and/or in rotations with herbicides with a different mode of action effective on the same weed species

RECOMMENDATION:

For the control of certain broad-leaved and grass weeds in all winter and spring sown cereals except oats.

Apply 150 g/ha Hussar with 1 litre/ha of Partner around the tillering stage of the crop but before, weeds reach the maximum growth stage specified. Do not apply to crops other than wheat, barley, ryecorn, triticale or durum wheat.

CROP SAFETY

Do not apply to crops under stress. Hussar can be used on all commercial cereal cultivars up to the 2nd node stage. Some minor temporary chlorosis and/or stunting may occur.

WEEDS CONTROLLED AT 150 G/HA + Partner at 1 litre/ ha.

Broad-leaved Weeds	Weed size	Grass Weeds	Weed size
Broad-leaved dock	Up to 8 leaf	Annual poa	Up to 2 tiller
Californian thistle	10-15 cm tall	Forage tall fescue	Up to 2 tiller
Chickweed	Up to 8 leaf	Grazing phalaris	Up to 2 tiller
Cleavers	2-3 whorl	Lesser canary grass	Up to 2 tiller
Fathen	Up to 8 leaf	Ryegrass-annual	Up to 2 tiller
Hawkesbeard	Up to 6 leaf	Ryegrass-hybrid	Up to 2 tiller
Hedge mustard	Up to 8 leaf	Ryegrass-perennial	Up to 2 tiller
Henbit	Up to 8 leaf		
Mouse-ear chickweed	Up to 6 leaf	Incomplete Control	
Nipplewort	Up to 6 leaf	Field pansy	



22-06-2018

Oxtongue	Up to 6 leaf	Parsley piert
Rayless chamomile	Up to 8 leaf	Speedwell
Red deadnettle	Up to 8 leaf	Staggerweed
Shepherd's purse	Up to 8 leaf	Browntop
Sow thistle	Up to 8 leaf	Cocksfoot
Spurrey	Up to 8 whorl	
Stinking mayweed	Up to 6 leaf	
Twin cress	Up to 8 leaf	

White clover	6 to 10 leaf	Not Controlled
Willow weed	Up to 6 leaf	ALS resistant Chickweed
Wireweed	Up to 6 leaf	Black nightshade
Yarrow	6 to 8 leaf	Brome Spp.
Yellow gromwell	Up to 8 leaf	

Best results are obtained from applications made to small actively growing weeds.

Weeds that are stressed may be suppressed only.

Some weeds emerging after application may not be controlled

COMPATABILITY: Hussar may be tank mixed with Cougar® to broaden the spectrum of broad-leaved weeds. However, DO NOT tank mix if the field to be sprayed contains target grass weeds as decreased efficacy will result.

RESTRICTIONS:

RAINFALL: Do not apply if rainfall is expected within 2 hours.

Undersown Crops: Do not apply to cereals undersown with grasses, clovers or other susceptible crops.

Crop Rotation Recommendations

In the event of a crop failure, allow the following intervals before sowing these crops:

Time - Months	Crops Safely Sown
0	Wheat, Barley, Maize
1	Peas, Beans, Oats
2	Ryegrass, Kale, Turnips
3	Clover and Pasture mixes

Thorough cultivation may allow these intervals to be reduced.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Commodities.

WITHHOLDING PERIOD:

Silage/Grazing: barley 28 days, other cereals (except barley) 56 days

Sprayer Clean Up

Failure to clean residues from spraying equipment can result in injury to sensitive crops.



22-06-2018

Ensure that the following operation is carried out in an area that is clear of waterways and desirable vegetation.

1. Drain sprayer, pump and lines completely and wash out with clean water.
2. Drain again.
3. Fill the tank with water and add 1 litre chlorine bleach (3% sodium hypochlorite) per 100 litres water with agitation running.
4. Flush some of the bleach solution through pump, lines and nozzles. Allow remainder to agitate in tank for 10 minutes.
5. Remove nozzles and filters and leave to soak in a bleach solution of 500 ml per 10 litres water while the tank cleaning is in progress.
6. Drain the tank and repeat the procedure of flushing with bleach solution.
7. Flush the tank and lines with clean water to remove traces of bleach.
8. Double rinse the nozzles and filters in clean water.

Dispose of surplus spray mix, rinseate and any spillage in accordance with the directions given in NZS 8409: 1999

CONDITIONS OF SUPPLY: READ THESE CONDITIONS OF SUPPLY BEFORE BUYING AND USING. IF THESE CONDITIONS ARE UNACCEPTABLE, RETURN AT ONCE UNOPENED.

To the extent permitted by law Bayer does not accept any liability or responsibility whatsoever for any loss, damage or injury whether direct, indirect or consequential howsoever arising in connection with these goods, including without limitation their storage, application, handling or use. Bayer does not make any warranties (express or implied) with respect to fitness for purpose, merchantable quality or any other matter.

To the extent only that Bayer is prevented by law from contracting out of the Consumer Guarantees Act, then these conditions apply subject to that Act. The Act shall not apply where these goods are acquired for business purposes.

Date of Manufacture:

Batch No:

Hussar[®], Cougar[®] and Partner[®] are registered trademarks of Bayer

Glean[®] is a registered trademark of E I Dupont de Nemours and Co Inc, Wilmington, Delaware 19898 USA

Bayer

Registered to:
Bayer New Zealand Limited
3 Argus Place, Hillcrest, Auckland 0627

Ph. 0800 428 246
EMERGENCY RESPONSE SERVICE (24 hrs) 0800 734 607



A handwritten signature in blue ink, appearing to be 'Mark Zimmmerman', written over the bottom right portion of the red stamp.